

TAISHO KAMPO LAXATIVES

- ◆ TAISHO KAMPO LAXATIVES contains the Chinese herbal medicine "Daiokanzoto". It is a laxative that relieves constipation and pimples, dermatitis and bloated stomach (abdominal distension) associated with constipation.
- ◆ Take the drug at bedtime, and you will have a natural bowel movement the following morning.
- ◆ The small tablets are scored and easy to take.

Indication

To alleviate such symptoms as constipation, dull headache/hot flush/eczema/dermatitis/pimples (acne)/loss of appetite/abdominal distention/abnormal colonic fermentation/hemorrhoid associated with constipation

Precaution

The drug is available regardless of your physical strength.

Dosage and administration

Take the following dose with cold or lukewarm water without chewing at bedtime.

However, start with a minimum dose and increase or decrease the dose by 1/2 or 1 tablet (up to the maximum dose below) depending on bowel movement or conditions.

Adults (15 years or over), when bowel movement is absent for 2 to 3 days: 1 to 3 tablets per dose, once daily

Adults (15 years or over), when bowel movement is absent for 4 days or more: 2 to 4 tablets per dose, once daily

Children (5 to 14 years), when bowel movement is absent for 2 to 3 days: 0.5 to 1.5 tablets per dose, once daily

Children (5 to 14 years), when bowel movement is absent for 4 days or more: 1 to 2 tablets per dose, once daily

Under 5 years: Do not take

Precaution

- Comply with the prescribed dosage and administration instructions.
- The use of the drug in children should be supervised by a parent.
- Women with a period should decrease the dose.
- Do not return the tablets touched by wet fingers into the jar. (It will cause discoloration and may deteriorate the quality)

Ingredient

In 4 tablets

Powdered Diokanzoto Extract 400mg

(Rhubarb 1600mg, glycyrrhiza 400mg)

Excipients

Magnesium aluminometasilicate, anhydrous silicic acid, potato starch, carboxymethylcellulose calcium, hydroxypropylcellulose, magnesium stearate, hardened oil

Precaution

- Do not worry if you have orange- or red-colored urine after taking the drug as this is caused by the Daio contained in the drug.
- The drug contains herbal medicine as a raw ingredient. The color of the drug may differ slightly, but this will not affect the efficacy.

Precautions

When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur)

- This medicine should not be taken together with the following medicines
Other cathartic agents
- For those who are lactating, do not use this medicine or avoid lactating while using this medicine

Consultation

- The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration
Patients undergoing medical treatment from a physician.
Pregnant women or women suspected of being pregnant.

Persons who is physically weak (declining of physical strength or constitutional weakness).

Persons with weak stomach and intestine who have a tendency to diarrhea, diarrhea with severe abdominal pain.

Patients who have experienced allergic symptoms associated with drugs, etc.

Persons with the following symptoms.

Severe abdominal pain, nausea and vomiting

- If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation

Skin: rash/redness, itching

Gastrointestinal system: diarrhea with severe abdominal pain, nausea and vomiting, abdominal pain

- The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation

Diarrhea

- When symptoms do not improve even after taking this drug for 5 to 6 days, stop taking this drug and consult a physician, pharmacist or registered salesperson, being sure to take this instruction leaflet with you

Precautions for storage and handling

- Store in a cool, low humidity place away from direct sunlight in an air-tight container.
- Store the product beyond the reach of children.
- Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)
- Do not use the product past the expiration date. Even before the expiration date, use the product within 6 months after it is opened. (to assure the quality)

21

【Disclaimer on Multilingual Product Information】

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.