

TAISHO STOMATITIS CHEWABLE TABLETS

Class 3 OTC drug
Stomatitis Products

◆ TAISHO STOMATITIS CHEWABLE TABLETS is an oral stomatitis drug that quickly dissolves in the mouth to exert a good effect on painful and smarting stomatitis. It can be taken without water and easily at any place.

Indication

stomatitis
Pharyngitis, tonsillitis (sore throat, and swollen throat.)

Dosage and administration

Take the following dose by chewing or dissolving the drug in the mouth.
Adults (15 years or over): 2 tablets per dose, 3 times daily
7 to 14 years: 1 tablet per dose, 3 times daily
Under 7 years: Do not take

ingredient and amount

In 6 tablets
Tranexamic acid
750mg
Dipotassium Glycyrrhizinate
63mg
Nicotinamide
60mg
Pyridoxine hydrochloride
50mg
Riboflavin
12mg

Excipients

cellulose, anhydrous silicic acid, Corn starch, D-Mannitol, Crospovidone, Aspartame (L- phenylalanine compound), Acesulfame K, Sucralose, Citric acid, Magnesium stearate, Flavoring, Starch sodium octenyl succinate

Precautions relating to ingredient and amount

No need to worry if you have yellow urine after taking the drug because it is caused by the vitamin B2 contained in the drug.

Precautions

When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur.)

- This medicine should not be taken together with any of the following medicines:
internal medicines containing glycyrrhiza or its main component glycolyl
internal medicines containing Tranexamic Acid (Oral Remedies for Rhinitis, cold medicine, Antipyretic Analgesics expectorants, etc.),
- Do not take this medicine for a long time

Consultation

- The following persons should contact a physician, dentist, pharmacist, or registered salesperson for a consultation before administration.

Pregnant women, nursing women, or women suspected of being pregnant.

The elderly

Persons who have had an allergic symptom to drugs.

Persons with the following symptoms:

oedema

Persons diagnosed as having the following

Hypertension

heart disease

kidney disease

Patients with blood clots (cerebral thrombosis, myocardial infarction, thrombophlebitis)

Patients who may develop thrombosis

- If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, dentist, pharmacist, or registered salesperson for a consultation.

Skin
rash/redness
itching
Gastrointestinal system
nausea and vomiting
Heartburn
Loss of appetite or increased appetite
Neuropsychiatric system
dizziness
Urinary system:
Frequent urination

The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid from a physician. Pseudoaldosteronism, myopathy: In addition to listlessness, numbness, stretched feeling, and stiffness of hands and feet, feeling of weakness and muscular pain will appear and gradually progress

● The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this document to your physician, dentist, pharmacist, or registered salesperson for a consultation.

diarrhea
● When symptoms do not improve even after taking the medicine 5 to 6 times, stop taking this medicine and consult a physician, dentist, pharmacist or registered salesperson, being sure to take this instruction leaflet with you.

Precautions for storage and handling

Avoid direct sunlight, and store the product in a cool place with little humidity.

Store the product beyond the reach of children.

Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration.)

After opening the inner bag (Aluminum bag), store it in a place where it will not be exposed to light, such as returning it to a box, in order to prevent discoloration due to light.

Do not use the product past the expiration date. Even before the expiration date, use the product within 6 months after it is opened (to assure the quality).

11

【Disclaimer on Multilingual OTC Product Information】

• This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.

• Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.

• Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.

• The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.