

STOMATITIS OINTMENT TAISHO A

Class 3 OTC drug
Stomatitis Products

◆STOMATITIS OINTMENT TAISHO A is an ointment developed to treat stomatitis and angular stomatitis. The ointment sticks well on the affected site, and covers the site even in the mouth for long hours.

Indication

Dry cracking lips/sores, stomatitis, and glossitis
angular stomatitis, canker sores

Dosage and administration

Apply an appropriate dose onto the lesion 2 to 4 times daily.
Comply with the prescribed dosage and administration instructions.
Clean the affected site and surrounding area before use.
Attention should be paid so that the medicine will not come into contact with eyes. In case the medicine comes into contact with eyes, immediately rinse them with water or lukewarm water, and immediately seek an ophthalmologist for medical treatment.
Use for children should be supervised by a parent.
Use the drug only for oral cavity and lips.
Do not use this drug with other external preparations on the same site.

ingredient and amount

In 100 g
Cetylpyridinium Chloride 0.1g
Allantoin 0.5g

Excipients

Xylitol ,Carboxyvinyl polymer,Hypromellose(Hydroxypropylmethylcellulose),Hydrocarbon gel,flavor,l-Menthol

Precautions

When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur.)
Do not use in the following sites.
Lesions with erosion

Consultation

- The following persons should contact a physician, dentist, pharmacist, or registered salesperson for a consultation before use.
Patients undergoing medical treatment from a physician or dentist.
infants and small children
Patients who have experienced allergic symptoms associated with drugs, etc.
- If the following symptoms are observed after using this drug, these may be adverse reactions, so immediately discontinue the use, and show this document to your physician, dentist, pharmacist, or registered sales person for a consultation.
Skin
the mouth
rash/redness
itching
- When symptoms do not improve even after using the medicine 5 to 6 days, stop using this medicine and consult a physician, dentist, pharmacist or registered salesperson, being sure to take this instruction leaflet with you.

Precautions for storage and handling

Store in a cool, low humidity place away from direct sunlight.
Store the product beyond the reach of children.
Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)
Do not use the product past the expiration date.
Even within the expiration date, use up as soon as possible once it is opened (to assure the quality).

【Disclaimer on Multilingual OTC Product Information】

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.