

LIPOVITAN fine

◆ Works when you are tired

Indication

- ★ Recovery from and prophylaxis of fatigue
- ★ Maintenance and improvement of physical strength or resistance or concentration
- ★ Improvement and prophylaxis of physical malfunction associated with malnutrition in daily living: getting tired easily, persisting fatigue, lack of physical strength, physical tiredness or feeling languid, bad skin conditions (rough or dry skin), eyestrain, bad conditions of the shoulders, neck, lower back or knees, decreased appetite and malaise associated with hangover
- ★ Improvement and prophylaxis of physical malfunction associated with frail constitution (including physical weakening due to aging): weakening of bones or teeth
- ★ Nutritional support for declined physical strength during/after illness, febrile debilitating illness, anorexia, pregnancy/lactation period, before and after childbirth, etc.

Dosage and administration

Adults (15 years or over): Take 1 bottle (100mL) per dose once daily.

Under 15 years: Do not use.

Precaution

Comply with the prescribed dosage and administration instructions. (Pay attention to excessive intake, etc. if a product containing other vitamins, etc. is taken together with this drug)

Ingredient

In 100mL

Taurine (aminoethyl sulfonic acid) 1000mg

Thiamine nitrate (vitamin B₁) 5mg

Riboflavin sodium phosphate (vitamin B₂) 10mg

Pyridoxine hydrochloride (vitamin B₆) 10mg

Nicotinamide 20mg

Carnitine hydrochloride 10mg

Anhydrous caffeine 50mg

Excipients

Erythritol, acesulfame K, purified stevia extract, sucralose, citric acid, sodium citrate, propyl gallate, sodium benzoate, flavoring, vitamin E, vanillin

Precaution

No need to worry if you have yellow urine after taking the drug because it is caused by the vitamin B₂ contained in the drug.

Precautions

Consultation

- If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this product to your physician, pharmacist, or registered salesperson for a consultation
 - Skin: rash
 - Gastrointestinal system: feeling of discomfort in the stomach
- The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this product to your physician, pharmacist, or registered salesperson for a consultation
 - Diarrhea
- When symptoms do not improve even after taking the medicine for a while, stop taking this medicine, and show this product to a physician, pharmacist, or registered salesperson for consultation

Precautions for storage and handling

- Store this in a cool place, avoiding direct sunlight.
- Store the product beyond the reach of children.
- Do not take the product past the expiration date.

21

【Disclaimer on Multilingual Product Information】

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.