

# LIPOVITAN D HYPER

Nutrients/tonics, Nutritional supplementation

## Indication

- ★ Nutritional supplements during physical fatigue, during and after illness, loss of appetite, nutritional disorders, feverish wearing disease, for nutritional supplementation during pregnancy/nursing
- ★ Nutrients/tonics
- ★ Weak constitution

## Dosage and administration

Adults (15 years or over): Take 1 bottle (100mL) per dose once daily.

Under 15 years: Do not use.

### Precaution

- Comply with the prescribed dosage and administration instructions. (Pay attention to excessive intake, etc. if a product containing other vitamins, etc. is taken together with this drug.)
- The drug contains a herbal extract. Sedimentation may occur in rare cases. This does not affect drug efficacy. Shake well before taking the drug.

## Ingredient

In 100mL

Taurine (aminoethyl sulfonic acid) 3000mg

Diisopropylamine dichloroacetate 20mg

Royal jelly 100mg

Carnitine hydrochloride 50mg

Thiamine nitrate (vitamin B<sub>1</sub>) 10mg

Riboflavin sodium phosphate (vitamin B<sub>2</sub>) 5mg

Pyridoxine hydrochloride (vitamin B<sub>6</sub>) 10mg

Inositol 100mg

Nicotinamide 20mg

Anhydrous caffeine 50mg

### Excipients

Sugar, D-sorbitol, acesulfame K, sucralose, purified stevia extract, citric acid, sodium citrate, povidone, propyl gallate, sodium benzoate, caramel, flavoring, glycerin

### Precaution

No need to worry if you have yellow urine after taking the drug because it is caused by the vitamin B<sub>2</sub> contained in the drug.

## Precautions

### Consultation

- In the following cases, immediately discontinue this drug, and consult your physician or pharmacist. Be sure to show this product  
If the following symptoms occur after taking the drug.
  - Skin: rash
  - Gastrointestinal system: feeling of discomfort in the stomachWhen symptoms do not improve even after taking the medicine for a while.
- The following symptoms may occur. If these symptoms persist or worsen, discontinue this drug, and consult your physician or pharmacist
  - Diarrhea

## Precautions for storage and handling

- Store this in a cool place, avoiding direct sunlight.
- Store the product beyond the reach of children.

【Disclaimer on Multilingual Product Information】

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.