

## LIPOVITAN DX AMINO

◆ LIPOVITAN DX AMINO contains amino acids such as BCAA\* (valine, leucine, and isoleucine) and arginine, as well as calcium, in addition to taurine and vitamins B<sub>1</sub>, B<sub>2</sub>, and B<sub>6</sub>, which are the base formulas of the LIPOVITAN series that has been helping people deal with their fatigue. A once-daily dosage is highly effective in relieving fatigue experienced in daily life.

\*Branched-chain amino acid

◆ It is also effective in improving and preventing muscle weakness associated with aging.

◆ This product is especially recommended for middle-aged and elderly people who feel easily fatigued and muscle weakness associated with aging, in order to stay vigorous and active every day.

### Indication

★ Recovery from and prophylaxis of fatigue

★ Improvement and prophylaxis of physical malfunction associated with frail constitution (including physical weakening due to aging)

Muscle weakness

Getting tired easily, persisting fatigue, lack of physical strength, physical tiredness or feeling languid

Bad conditions of the shoulders, neck, lower back or knees

weakening of bones or teeth

★ Maintenance and improvement of physical strength or resistance or concentration

★ Improvement and prophylaxis of physical malfunction associated malnutrition in daily living

Muscle weakness

Getting tired easily, persisting fatigue, lack of physical strength, physical tiredness or feeling languid

Bad conditions of the shoulders, neck, lower back or knees

★ Nutritional support for declined physical strength during/after illness, febrile debilitating illness, anorexia, pregnancy/lactation period, before and after childbirth, etc.

### Dosage and administration

Take the following dose with cold or lukewarm water.

Adults (15 years or over): 3 tablets per dose, once daily

Under 15 years: Do not use

### Precaution

- Comply with the prescribed dosage and administration instructions. (Pay attention to excessive intake, etc. if a product containing other vitamins, etc. is taken together with this drug)
- Do not return the tablets touched by wet fingers into the jar. (This causes discoloration and may deteriorate tablet quality)
- The small package wrapped with paper in the jar is a drying agent. Do not take it.

### Ingredient

In 3 tablets

Taurine (aminoethyl sulfonic acid) 200mg

Thiamine nitrate (vitamin B<sub>1</sub>) 10mg

Riboflavin (vitamin B<sub>2</sub>) 5mg

Pyridoxine hydrochloride (vitamin B<sub>6</sub>) 5mg

L-Valine 50mg

L-Leucine 100mg

L-Isoleucine 50mg

L-Arginine hydrochloride 30mg

Magnesium L-aspartate 180mg

L-histidine hydrochloride hydrate 6mg

Anhydrous dibasic calcium phosphate 102mg (30mg as calcium)

## Excipients

Anhydrous silicic acid, cellulose, hydroxypropylcellulose, magnesium stearate, talc, macrogol, titanium oxide, hypromellose, ferric oxide, carnauba wax

## Precaution

No need to worry if you have yellow urine after taking the drug because it is caused by the vitamin B<sub>2</sub> contained in the drug.

## Precautions

### Consultation

- If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation
  - Skin: rash
  - Gastrointestinal system: feeling of discomfort in the stomach
- The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation
  - Diarrhea
- When symptoms do not improve even after taking this medicine for a while, stop taking this medicine and consult a physician, pharmacist or registered salesperson, being sure to take this instruction leaflet with you

## Precautions for storage and handling

- Store in a cool, low humidity place away from direct sunlight.
- Store the product beyond the reach of children.
- Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)
- Do not use the product past the expiration date. Even before the expiration date, use the product within 6 months after it is opened. (to assure the quality)

11

### 【Disclaimer on Multilingual Product Information】

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.