

TAISHO ANTIDIARRHEALS <for PEDIATRIC USE>

Class 2 OTC drug
Antidiarrheals

◆ TAISHO ANTIDIARRHEALS (for PEDIATRIC USE) relieves uncomfortable diarrhea symptoms, and supplements vitamins B 1 and B 2 which tend to be consumed by diarrhea. ■ The banana-flavored drug is easy for little children to take.

Indication

Diarrhea, diarrhea due to dyspepsia, food poisoning, vomiting and diarrhea, water poisoning, diarrhea, and soft stools

Dosage and administration

Take the following dose with cold or lukewarm water up to 3 times daily. Provide a dosing interval for at least 4 hours.

5 to 7 years: 1 packet per dose, up to 3 times daily

3 to 4 years: 2/3 packet per dose, up to 3 times daily

1 to 2 years: 1/2 packet per dose, up to 3 times daily

3 months to under 1 year: 1/4 packet per dose, up to 3 times daily

Under 3 months: Do not take

Comply with the prescribed dosage and administration instructions.

The use of the drug in children should be supervised by a parent.

For infants under 1 year old, treatment by a physician should be prioritized and, when only necessary, allow them to take the medicine.

ingredient and amount

In 1 packet (1.2 g)

Berberine Tannate 30 mg 30mg

Albumin Tannate 440mg

Biodiastase2000 30mg

Thiamine Nitrate 2mg

Riboflavin 1mg

Excipients

Potato starch, Sugar, Corn starch, Magnesium aluminometasilicate, Hydroxypropylcellulose, flavor, Gelatin, Sorbitan fatty acid ester

Precautions relating to ingredient and amount

No need to worry if you have yellow urine after taking the drug because it is caused by the vitamin B2 contained in the drug.

Precautions

When not to use the product

Although the drug is intended for children, the document describes the general precaution for antidiarrheal (If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur.)

● This product should not be taken in the following persons.

Patients who have had an allergic symptom to this drug, its ingredients, or cow milk

Consultation

● The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.

Patients undergoing medical treatment from a physician

Those who have diarrhea with fever, bloody stool or persistent mucous stool.

Those who have acute and severe diarrhea or diarrhea with accompanying symptoms such as abdominal pain/abdominal distension/sick feeling. (The disease may worsen by taking this medicine to stop diarrhea.)

The elderly

Patients who have experienced allergic symptoms associated with drugs, etc.

● If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

Skin

rash/redness

itching

The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid from a physician.

Shock (anaphylaxis): Symptoms, such as itching of skin, urticaria, hoarseness, sneezing, itchy throat, breathing difficulties, palpitations, and clouding of consciousness may occur immediately after taking the drug.

If symptoms do not resolve after using the drug for 5 or 6 days, stop using this medicine, and show this document to your

physician, pharmacist, or registered salesperson for a consultation.

Precautions for storage and handling

Avoid direct sunlight, and store the product in a cool place with little humidity.

Store the product beyond the reach of children.

Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)

If you divide a pack to take the rest afterward, fold up the opening of the bag to store, and take it within 2 days.

Do not take the product past the expiration date.

72

【Disclaimer on Multilingual OTC Product Information】

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.