

## PABRON GOLD A <TABLETS>

- ◆ PABRON GOLD A <TABLETS> is a cold medication that contains 7 active ingredients such as guaifenesin, and relieves symptoms associated with the common cold, such as cough, sputum, and sore throat.
- ◆ It is suitable for family use.

### Indication

To alleviate cold symptoms (cough, sputum, sore throat, sneezing, runny nose, stuffy nose, chills, fever, headache, joint pain, and muscle pain)

### Dosage and administration

Take the following dose with cold or lukewarm water within 30 minutes after a meal whenever possible.

15 years or over: 3 tablets per dose, 3 times daily

12 to 14 years: 2 tablets per dose, 3 times daily

Under 12 years: Do not take

### Precaution

- Comply with the prescribed dosage and administration instructions.
- The use of the drug in children should be supervised by a parent.
- Do not return the tablets touched by wet fingers into the jar. (It will cause discoloration and may deteriorate the quality)

### Ingredient

In 3 tablets

Guaifenesin 60mg

Dihydrocodeine Phosphate 8mg

dl-Methylephedrine Hydrochloride 20mg

Acetaminophen 300mg

Chlorpheniramine Maleate 2.5mg

Anhydrous caffeine 25mg

Riboflavin (vitamin B<sub>2</sub>) 4mg

### Excipients

Cellulose, anhydrous silicic acid, magnesium aluminometasilicate, hydroxypropylcellulose, sodium starch glycolate, lactose, magnesium stearate, hardened oil, flavoring, vanillin, benzyl benzoate

### Precaution

No need to worry if you have yellow urine after taking the drug because it is caused by the vitamin B<sub>2</sub> contained in the drug.

### Precautions

#### When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur)

- This product should not be taken in the following persons
  - Patients who have had an allergic symptom to this drug or its ingredients.
  - Patients who have experienced asthma from taking this drug or other cold remedies, antipyretic analgesics.
  - Children under 12 years old.
- Do not use any of the following medications while taking this drug
  - Other cold medicines, antipyretics/analgesics, sedatives, expectorants, and internal medicines, etc. containing antihistamines (internal medicine for rhinitis, medicine for motion sickness, medicine for allergies, etc.)
- After taking this drug, do not drive a car or operate machinery (symptoms such as sleepiness may occur)
- For those who are lactating, do not use this medicine or avoid lactating while using this medicine
- Do not drink alcohol before and after taking this drug
- Do not take this medicine for a long time

## Consultation

- The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration
  - Patients undergoing medical treatment from a physician or dentist.
  - Pregnant women or women suspected of being pregnant.
  - The elderly.
  - Patients who have experienced allergic symptoms associated with drugs, etc.
  - Persons with the following symptoms.
    - Hyperthermia, urination difficulty
  - Persons diagnosed as having the following.
    - Thyroid disease, diabetes, heart disease, hypertension, liver disease, kidney disease, gastric/duodenal ulcer, glaucoma, respiratory functional disorder, obstructive sleep apnea syndrome, adiposity
- If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation
  - Skin: rash/redness, itching
  - Gastrointestinal system: nausea/vomiting, and loss of appetite
  - Neuropsychiatric system: dizziness
  - Urinary system: urination difficulty
  - Other: excessive temperature decrease
- The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid.
  - Shock (anaphylaxis): Symptoms, such as itching of skin, urticaria, hoarseness, sneezing, itchy throat, breathing difficulties, palpitations, and clouding of consciousness may occur immediately after take.
  - Oculomuocutaneous syndrome (stevens-johnson syndrome), toxic epidermal necrolysis, acute generalised exanthematous pustulosis: hyperthermia, ocular hyperaemia, eye discharge, lip erosion, pain throat, widespread skin rash/redness, small pimples (small pustules) on reddened skin, general malaise, anorexia, etc. may persist or suddenly worsen.
  - Drug-induced hypersensitivity syndrome: Symptoms such as widespread redness of the skin, generalised rash, fever, malaise, and swelling of lymph nodes (neck, armpit, groin, etc.) may occur.
  - Hepatic function failure: Symptoms, such as fever, itching, rash, jaundice (yellowing of skin and white of eyes), brown urine, general malaise, loss of appetite, etc. may occur.
  - Kidney disorders: Symptoms, such as fever, rash, reduced urinary volume, general oedema, general malaise, arthralgia (painful joints) and diarrhea, etc. may occur.
  - Interstitial pneumonia: Shortness and/or difficulties of breath when go upstairs or overwork, sudden dry cough and/or fever and its continuance.
  - Asthma: Symptoms, such as wheezing or hissing when breathing, and difficult breathing, etc. may occur.
  - Aplastic anemia: symptoms, such as bruises, nose bleeds, bleeding gums, fever, pale appearance of skin and mucosa, fatigue, palpitations, shortness of breath, becoming dizzy from feeling sick and blood in the urine, etc. may occur.
  - Agranulocytosis: Symptoms, such as acute high fever, chills and sore throat, etc. may occur.
  - Respiratory depression: shortness of breath, breathing difficulties, etc. may occur.
- The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation
  - Constipation, dry mouth, drowsiness
- When symptoms do not improve even after taking the medicine 5 to 6 times, stop taking this medicine and consult a physician, pharmacist or registered salesperson, being sure to take this instruction leaflet with you

## Precautions for storage and handling

- Store in a cool, low humidity place away from direct sunlight in an air-tight container.
- Store the product beyond the reach of children.
- Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)
- Do not use the product past the expiration date. Even before the expiration date, use the product within 6 months after it is opened. (to assure the quality)

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.