

Make sure to read this instruction sheet before taking this medicine.
Additionally, store this sheet so that it can be referenced when needed.

Take this medicine twice a day for severe runny nose, stuffy nose,
and sneezing <long-acting rhinitis medicine>

PABRON RHINITIS CAPSULES Sa

Brand name: PABRON RHINITIS CAPSULES Sa Class 2 OTC drug <Rhinitis medicine>

- ◆ PABRON RHINITIS CAPSULES Sa is an oral rhinitis medicine that is highly effective when taken twice a day.
- ◆ The white granules will dissolve and start working quickly, while the orange granules will dissolve and start working slowly to effectively relieve severe symptoms of rhinitis.



Precautions for Use



Things You Must Not Do

(Failure to follow these precautions may lead to your current symptoms worsening or a greater chance of side effects or accidents occurring)

1 The following individuals should not take this medicine:

- (1) Persons who have had allergic symptoms as a result of taking this medicine or any component of it.
- (2) Persons with the following symptoms:
Difficulty urinating due to enlarged prostate
- (3) Persons who have received any of the following diagnoses:
Hypertension, heart disease, thyroid dysfunction, diabetes

2 Do not use any of the following medicines while taking this medicine:

Other oral medicines for rhinitis, oral medicines containing antihistamines, etc. (cold medicines, antitussive expectorants, motion sickness medicines, allergy medicines, etc.), oral medicines containing pseudoephedrine hydrochloride or pseudoephedrine sulfate, gastrointestinal analgesic/antispasmodic medicines

3 Do not drive a vehicle or operate machinery after taking this medicine
(You may experience symptoms such as drowsiness, blurred vision, abnormal glare, etc.)

4 Do not use for long periods of time



Things To Consult On

1 The following individuals should consult with their doctor, pharmacist, or a registered seller before taking this medicine

- (1) Persons who are currently being treated by a doctor.
- (2) Persons who are pregnant or who think they may be pregnant.
- (3) Persons who are breastfeeding.
- (4) Elderly persons.
- (5) Persons who have had allergic symptoms due to medication in the past.
- (6) Persons who have experienced insomnia, dizziness, weakness, tremors, or palpitations while taking cold medicines, antitussive expectorants, oral medicines for rhinitis, etc.
- (7) Persons with the following symptoms:
High fever, difficulty urinating
- (8) Persons who have received any of the following diagnoses:
Glaucoma, kidney disease
- (9) Persons receiving treatment with monoamine oxidase inhibitors (selegiline hydrochloride, etc.).
(Selegiline hydrochloride is used for treatment of Parkinson's disease)

2 If any of the following symptoms appear after you have taken this medicine, you may be experiencing side effects. As such, discontinue use immediately and consult with your doctor, pharmacist or registered seller, bringing this instruction sheet with you

Relevant Area	Symptoms
Skin	Rash, redness, itching
Digestive system	Nausea/vomiting, loss of appetite
Neuropsychiatric system	Dizziness, insomnia, nervousness, headache, convulsion
Urinary tract	Difficulty urinating
Other	Facial flushing, abnormal glare

In rare cases, the following serious symptoms may occur. In such cases, see a doctor immediately.

Symptom Name	Symptoms
Acute generalized exanthematous pustulosis	High fever, widespread skin rash/redness, small bumps (small pustules) appearing on reddened skin, whole body sluggishness and loss of appetite, etc., that persists or worsens rapidly

3 The following symptoms may appear after taking this medicine. If these symptoms persist or worsen, discontinue use and consult with your doctor, pharmacist, or registered seller, bringing a copy of this instruction sheet with you
Dry mouth, drowsiness, constipation, blurred vision

4 If symptoms do not improve after using this medicine for 5 to 6 days, discontinue use and consult with your doctor, pharmacist or registered seller, bringing this instruction sheet with you

Efficacy


Relieves the following acute rhinitis, allergic rhinitis, or sinusitis symptoms:

Sneezing, runny nose (excessive nasal discharge), stuffy nose, teary eyes, sore throat, dull headache (head feels heavy)



Usage/Dosage

Take the following dosage every 12 hours with room-temperature or lukewarm water.

Age	Single dose	Number of doses
15 years of age or older	2 capsule 	Twice per day
Under 15 years old	Do not take this medicine	

Note:

- (1) Strictly observe the prescribed dosage and administration.
- (2) How to take out the capsules

As shown in the illustration, press the convex part of the PTP sheet containing the capsule hard with a fingertip to tear the aluminum foil on the reverse side, remove the capsule, and take it. (Accidental ingestion of the PTP sheet as-is may lead to unexpected accidents, such as penetration into the esophageal mucosa.)

<Capsule removal diagram>



Ingredients

In 2 capsule

Ingredients	Amount	Action
Pseudoephedrine hydrochloride	60 mg	Vasoconstricting effect suppresses hyperemia/swelling of the mucous membranes of the nose and improves stuffy nose.
Carbinoxamine maleate	6 mg	Antihistamine effect relieves sneezing, runny nose, and stuffy nose.
Belladonna total alkaloid	0.2 mg	Secretion-inhibiting effect relieves runny nose and improves teary eyes.
Anhydrous caffeine	50 mg	Relieves dull headaches associated with rhinitis.

Excipients: magnesium aluminometasilicate, white sugar, corn starch, hydroxypropyl cellulose, magnesium stearate, cellulose, talc, ammonioalkyl methacrylate copolymer, stearyl alcohol, sorbitan trioleate, Yellow 5, gelatin, sodium lauryl sulfate

Precautions for Storage and Handling

- (1) Store in a cool, dry place away from direct sunlight.
- (2) Keep out of reach of children.
- (3) Do not replace the medicine into a different container. (This may result in misuse or a change in quality)
- (4) Do not take the product after the expiration date. Use this medicine within 6 months of opening, even if still within the expiration date. (In order to ensure quality.)



Disclaimer on Multilingual OTC Product Information

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.