

PABRON NASAL SPRAY

Class 2 OTC drug

nasal drop

◆PABRON NASAL SPRAY releases misty particles of active ingredients evenly deep in the nose, where active ingredients quickly suppress inflammation in nasal mucosa, and relieve stuffy nose. The capped container is convenient for carrying around.

Indication

To alleviate the following symptoms of acute rhinitis, allergic rhinitis or sinusitis: stuffy nose, runny nose (excessive nasal discharge), sneezing, and dull headache

Dosage and administration

Spray once to twice per dose into the nasal cavity. You may use the drug up to 6 times daily at an interval of at least 3 hours.

Adults (15 years or over) and children aged 7 years or over

Spray once to twice per dose into the nasal cavity.

Up to 6 times per day.

Under 7 years: Do not use

Comply with the prescribed dosage and administration instructions.

Excessive use may cause stuffy nose.

Use for children should be supervised by a parent.

Use only as a nose drop.

ingredient and amount

In 100 mL

Naphazoline Hydrochloride 0.05g

Chlorpheniramine Maleate 0.5g

Benzethonium chloride 0.02g

Excipients

Glycerin

Precautions

When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur.)

- Do not take this medicine for a long time

Consultation

- The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before use.

Patients undergoing medical treatment from a physician

Pregnant women or women suspected of being pregnant.

Patients who have experienced allergic symptoms associated with drugs, etc.

Persons diagnosed as having the following

Hypertension

heart disease

Diabetes

thyroid disease

Glaucoma

If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

Skin

rash/redness

itching

Nose:

swelling

Irritation

If symptoms do not resolve after using the drug for about 3 days, stop taking this medicine, and contact a physician, pharmacist, or registered salesperson for a consultation.

Precautions for storage and handling

Avoid direct sunlight, and store the product in a cool place with an air tight closure on.

Store the product beyond the reach of children.

Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)

Do not share the drug with other people.

Do not take the product past the expiration date.

【Disclaimer on Multilingual OTC Product Information】

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.