

PABRON KIDS COLD SYRUP

Class 2 OTC drug

Cold Remedies

◆PABRON KIDS COLD SYRUP is an oral cold medication for children from 3 months old. ■The strawberry-flavored syrup is easy to drink, and relieves such symptoms as cough, runny nose, and fever. ■The product does not contain caffeine or dl-Methylephedrine hydrochloride that disturb sleep.

Indication

To alleviate cold symptoms (sore throat, fever, cough, sputum, runny nose, stuffy nose, sneezing, chill (chill due to fever), headache, joint pain, muscle pain)

Dosage and administration

Measure the following dose with the graduated cup enclosed, and take the drug after a meal, and at bedtime if necessary, 3 times daily. In some cases, you may take the drug up to 6 times daily at an interval of about 4 hours.

3 to 6 years: 10 mL per dose, 3 times daily

1 to 2 years: 7.5 mL per dose, 3 times daily

6 months to under 1 year: 6 mL per dose, 3 times daily

3 months to under 6 months: 5 mL per dose, 3 times daily

Under 3 months: Do not take

Comply with the prescribed dosage and administration instructions.

The use of the drug in children should be supervised by a parent.

Infants under 2 years old should seek medical care from a physician first. Use the drug only in an unavoidable case.

Wipe the mouth of the jar with a clean gauze before closing the cap. (The solidified liquid may make the cap difficult to open.)

ingredient and amount

In 60 mL

Dextromethorphan Hydrobromide 16mg

Guaifenesin 83.3mg

Chlorpheniramine Maleate 2.5mg

Acetaminophen 300mg

Excipients

Sugar,Sucralose ,Benzoic acid,Butyl parahydroxybenzoate,Sodium citrate,citric acid,Sunset Yellow FCF ,flavor,Vanillin

Precautions

When not to use the product

Although the drug is intended for children, the document describes the general precaution for cold medicine (If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur.)

This product should not be used in the following persons:

Patients who have had an allergic symptom to this drug or its ingredients.

Patients who have experienced asthma from taking this drug or other cold remedies, antipyretic analgesics.

● This medicine should not be taken together with any of the following medicines:

Other cold medicines, antipyretics/analgesics, sedatives, expectorants, internal medicine, etc. containing antihistamines (internal medicine for rhinitis, medicine for motion sickness, medicine for allergies, etc.) and gastrointestinal analgesics

● After taking this drug, do not drive a car or operate machinery (sleepiness may occur).

Do not drink alcohol before and after taking this drug.

● Do not take this medicine for a long time

Consultation

● The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.

Patients undergoing medical treatment from a physician or dentist.

Pregnant women or women suspected of being pregnant.

Patients who have experienced allergic symptoms associated with drugs, etc.

Persons with the following symptoms:

hyperthermia

urination difficulty

Persons diagnosed as having the following

heart disease

Liver disease

kidney disease

Gastric/duodenal ulcer

Glaucoma

● If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

Skin

rash/redness

itching

Gastrointestinal system

Vomiting

Vomiting

Loss of appetite

Neuropsychiatric system

dizziness

Respiratory system:

shortness of breath

breathing difficulties/difficulty in breathing

Urinary system:

urination difficulty

Other:

Excessive temperature decrease

The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid from a physician.

Shock (anaphylaxis): Symptoms, such as itching of skin, urticaria, hoarseness, sneezing, itchy throat, breathing difficulties, palpitations, and clouding of consciousness may occur immediately after take.

Oculomucocutaneous syndrome (Stevens-Johnson syndrome), toxic epidermal necrolysis, acute generalised exanthematous pustulosis: Hyperthermia, ocular hyperaemia, eye discharge, lip erosion, pain throat, widespread skin rash/redness, small pimples (small pustules) on reddened skin, general malaise, anorexia, etc. may persist or suddenly worsen.

Hepatic function failure: Symptoms, such as fever, itching, rash, jaundice (yellowing of skin and white of eyes), brown urine, general malaise, loss of appetite, etc. may occur.

Kidney disorders: Symptoms, such as fever, rash, reduced urinary volume,...

Precautions for storage and handling

Avoid direct sunlight, and store the product in a cool place with an air tight closure on.

Store the product beyond the reach of children.

Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)

Do not take the product past the expiration date.

Even before the expiration date, use up to within 6 months after it is opened (to assure quality).

21

【Disclaimer on Multilingual OTC Product Information】

• This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.

• Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.

• Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.

• The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.