

PABRON SUPPOSITORIES for KIDS

- ◆ Fever is the most common symptom as it is said that all the pediatric diseases start with fever. However, it is important to alleviate fever because it could consume physical strength.
- ◆ PABRON SUPPOSITORIES for KIDS effectively relieves acute fever in children.

Indication

To temporarily alleviate pediatric fever

Dosage and administration

Insert the following dose into the anus.

6 to 12 years: 1 to 2 pieces per dose, once daily

3 to 5 years: 1 piece per dose, once daily

1 to 2 years: 1/2 to 1 piece per dose, once daily

Under 1 year: Do not use

Precaution

- Comply with the prescribed dosage and administration instructions.
- When allowing children to use the medicine, a guardian must be present to watch and tell such them how to use it.
- Do not use the drug for 2 days in a row.
- Use the drug only for the anus. Do not take it orally.

Ingredient

In 1 piece (1g)

Acetaminophen 100mg

Excipients

Hard fat

Precautions

Although the drug is intended for children, the document describes the general precaution for antipyretics.

When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur)

- This product should not be used in the following persons
 - Patients who have had an allergic symptom to this drug or its ingredients.
 - Patients who have experienced asthma from using this drug, other antipyretic analgesics, or cold remedies.
- Do not use any of the following medications while using this drug
 - Other antipyretics/analgesics, cold medicine, sedatives
- Do not drink alcohol before or after using this drug
- Do not take this medicine for a long time

Consultation

- The following persons should contact a physician, dentist, pharmacist, or registered salesperson for a consultation before use
 - Patients undergoing medical treatment from a physician or dentist.
 - Pregnant women or women suspected of being pregnant.
 - Elderly or weak persons.
 - Patients who have experienced allergic symptoms associated with drugs, etc.
 - Persons diagnosed as having the following.
 - Heart disease, kidney disease, liver disease, gastric/duodenal ulcer
- If the following symptoms are observed after using this drug, these may be adverse reactions, so immediately discontinue the use, and show this document to your physician, pharmacist, or registered sales person for a consultation
 - Skin: rash/redness, itching
 - Gastrointestinal system: nausea/vomiting, and loss of appetite, diarrhea, soft stool, urge to defecate

Neuropsychiatric system: dizziness

Other: excessive temperature decrease

- The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid.

Shock (anaphylaxis): Symptoms, such as itching of skin, urticaria, hoarseness, sneezing, itchy throat, breathing difficulties, palpitations, and clouding of consciousness may occur immediately after use.

Oculomucocutaneous syndrome (stevens-johnson syndrome), toxic epidermal necrolysis, acute generalised exanthematous pustulosis: hyperthermia, ocular hyperaemia, eye discharge, lip erosion, pain throat, widespread skin rash/redness, small pimples (small pustules) on reddened skin, general malaise, anorexia, etc. may persist or suddenly worsen.

Drug-induced hypersensitivity syndrome: Symptoms such as widespread redness of the skin, generalised rash, fever, malaise, and swelling of lymph nodes (neck, armpit, groin, etc.) may occur.

Hepatic function failure: Symptoms, such as fever, itching, rash, jaundice (yellowing of skin and white of eyes), brown urine, general malaise, loss of appetite, etc. may occur.

Kidney disorders: Symptoms, such as fever, rash, reduced urinary volume, general oedema, general malaise, arthralgia (painful joints) and diarrhea, etc. may occur.

Interstitial pneumonia: Shortness and/or difficulties of breath when go upstairs or overwork, sudden dry cough and/or fever and its continuance.

Asthma: Symptoms, such as wheezing or hissing when breathing, and difficult breathing, etc. may occur.

- If fever recurs after taking this drug once, stop taking this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

Precautions for storage and handling

- Avoid direct sunlight, and store in a cool place with little humidity and temperature below 30°C.
- Store the product beyond the reach of children.
- Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)
- Place the suppositories in the outer box with their tips downward, and store them upright as indicated on the outer box.
- Do not use the product past the expiration date. Even within the expiration date, try to use it up as soon as possible once it is opened. (to assure the quality)

11

【Disclaimer on Multilingual Product Information】

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.