Newly designated quasidrugs

Health drinks

Indication

- ☆ Maintenance and improvement of physical strength or resistance or concentration
- ☆ Recovery from and prophylaxis of fatigue
- ☆ Improvement and prophylaxis of physical malfunction associated malnutrition in daily living:Getting tired easily, persisting fatigue, lack of physical strength, physical tiredness or feeling languid
- ☆ Nutritional support for declined physical strength during/after illness, febrile debilitating illness, anorexia, pregnancy/lactation period, before and after childbirth, etc.

Dosage and administration

Adults (15 years or over): Take 1 bottle (50 mL) per dose once daily.

Under 15 years: Do not take

[Precaution]

Comply with the prescribed dosage and administration instructions.

(Pay attention to excessive intake, etc. if a product containing other vitamins, etc. is taken together with this drug.) The drug contains a herbal extract. Sedimentation may occur in rare cases. This does not affect drug efficacy. Shake well before taking the drug.

ingredient and amount

In 1 bottle (50 mL)

Ginger extract 77.8mg(equivalent to 700 mg of the crude drug)

Ginger Fluid Extract 0.3mL(equivalent to 300 mg of the crude drug)

Cinnamon Bark Fluid Extract 0.15mL(equivalent to 150 mg of the crude drug)

Peony Root extract 30mg(equivalent to 120 mg of the crude drug)

Jujube Extract 300mg(equivalent to 750 mg of the crude drug)

Glycyrrhiza Extract 30mg(equivalent to 120 mg of the crude drug)

Japanese Angelica Root Fluid ExtractS 0.3mL(equivalent to 300 mg of the crude drug)

Citrus Unshiu Peel Extract 20mg(equivalent to 100 mg of the crude drug)

Ginseng extract-P 85.8mg (equivalent to 600 mg of the crude drug)

Acanthopanax Fluid Extract 0.2mL(equivalent to 200 mg of the crude drug)

Dioscorea Rhizome Fluid Extract-A 0.2mL(equivalent to 200 mg of the crude drug)

Taurine 500mg

Riboflavin sodium phosphate 12mg

Pyridoxine hydrochloride (vitamin B6) 10mg

Nicotinamide 20mg

Excipients

Sugar, citric acid, Sodium citrate, Polyoxyethylene hydrogenated castor oil, Povidone, Sodium benzoate, Butyl parahydroxybenzoate, Caramel, flavor, Alcohol (not more than 0.39 mL)

[Precaution]

No need to worry if you have yellow urine after taking the drug because it is caused by the vitamin B2 contained in the drug.

Precautions

Consultation

• If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this product to your physician, pharmacist, or registered salesperson for a consultation.

Skin: rash

Gastrointestinal system: feeling of discomfort in the stomach

- The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this product to your physician, pharmacist, or registered salesperson for a consultation. Diarrhea
- When symptoms do not improve even after taking the medicine for a while, stop taking this medicine, and show this product to a physician, pharmacist, or registered salesperson for consultation.

Precautions for storage and handling

Store this in a cool place, avoiding direct sunlight.

14-Sep-2020 1/2 page

21

[Disclaimer on Multilingual OTC Product Information]

- •This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- •Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- •Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- •The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.

14-Sep-2020 2/2 page