

SEMPER RAMCURE

Class 2 OTC drug
Motion Sickness Products

◆ SEMPER RAMCURE is a medication for motion sickness that can be chewed and taken without water when you are busy before going out or feel sick during trip. ◆ It a grape-flavored tablet that children like and can be easily drunk like lemonade.

Indication

1) prevention and relief of dizziness, nausea, and headache associated with motion sickness

Dosage and administration

Take the following dose by chewing or dissolving the drug in the mouth.

To prevent motion sickness, take the drug 30 minutes prior to boarding.

If you need to take an additional dose, then take 1 dose of the drug at an interval of at least 4 hours.

11 years or over: 2 tablets per dose, up to twice daily

5 to 10 years: 1 tablet per dose, up to twice daily

Under 5 years: Do not use

Comply with the prescribed dosage and administration instructions.

When allowing children to take the medicine, a guardian must be present to watch and tell such them how to take it.

How to take out the tablets

•As shown in the figure, press the convex part of the PTP sheet containing the tablet firmly with your fingertips, break the aluminum foil on the back side, and take out the tablets.

• (If the tablets are inadvertently taken together with the PTP sheet containing them, it will lead to unexpected accidents, such as the sharp edge sticking into the esophageal mucosa.)

ingredient and amount

In 2 tablets

d-Chlorpheniramine Maleate

1.32mg

Scopolamine Hydrobromide

0.16mg

Excipients

Reduced maltose syrup, D-Mannitol , Potato starch, anhydrous silicic acid , Hydroxypropylcellulose, Aspartame (L-phenylalanine compound), Sucralose , Magnesium stearate, Ferric oxide , Flavoring

Precautions

When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur.)

●This drug should not be taken together with the following drugs:

Other medicine for motion sickness, cold medicine, antipyretics/analgesics, sedatives, expectorant, indigestion and heartburn relief, internal medicine containing antihistamines, etc. (such as internal medicine for rhinitis and medicine for allergies)

●After taking this drug, do not drive a car or operate machinery (symptoms such as sleepiness, blurred vision and abnormal brightness may occur).

Consultation

●The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.

Patients undergoing medical treatment from a physician

Pregnant women or women suspected of being pregnant.

The elderly

Persons who have had an allergic symptom to drugs.

Persons with the following symptoms:

urination difficulty

Persons diagnosed as having the following

Glaucoma

heart disease

Skin

rash/redness

itching

Neuropsychiatric system

headache

Urinary system:

urination difficulty

Other:

Facial hot flashes

abnormal brightness

The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid from a physician.

Aplastic anemia: Symptoms, such as bruises, nose bleeds, bleeding gums, fever, pale appearance of skin and mucosa, fatigue, palpitations, shortness of breath, becoming dizzy from feeling sick and blood in the urine, etc. may occur.

Agranulocytosis: Symptoms, such as acute high fever, chills and sore throat, etc. may occur.

●The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.

Thirst, sleepiness, constipation, blurred vision

Precautions for storage and handling

Avoid direct sunlight, and store the product in a cool place with little humidity.

Store the product beyond the reach of children.

Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration.)

Do not use the product past the expiration date. Even within the expiration date, try to use it up as soon as possible once it is opened (to assure the quality).

11

【Disclaimer on Multilingual OTC Product Information】

•This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.

•Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.

•Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.

•The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.