

SEMPER PRO

- ◆ SEMPER PRO prevents and alleviates symptoms of severe motion sickness such as dizziness, nausea, and headache.
- ◆ You can use the drug if you are going to travel by car, bus, airplane, or boat that rolls a lot, or to an amusement park ride.
- ◆ Once daily oral dose takes effect.
- ◆ The drug relieves symptoms when you take it even after you begin to feel sick.

Indication

Prevention and relief of dizziness, nausea, and headache associated with motion sickness

Dosage and administration

Take the following dose with cold or lukewarm water.

However, to prevent motion sickness, take the drug 30 minutes prior to boarding.

15 years or over: 1 tablets per dose, 1 times daily

Under 15 years: Do not take

Precaution

Comply with the prescribed dosage and administration instructions.

How to take out the tablets

Press the convex part of the PTP sheet containing the tablet firmly with your fingertips, break the aluminum foil on the back side, and take out the tablets. (If the tablets are inadvertently taken together with the PTP sheet containing them, it will lead to unexpected accidents, such as the sharp edge sticking into the esophageal mucosa)

Ingredient

In 1 tablets

Ethyl aminobenzoate 100mg

Scopolamine Hydrobromide 0.25mg

d-Chlorpheniramine Maleate 2mg

Pyridoxine hydrochloride (vitamin B₆) 10mg

Anhydrous caffeine 20mg

Excipients

Anhydrous silicic acid, cellulose, D-mannitol, hydroxypropylcellulose, sodium starch glycolate, magnesium stearate, ferric oxide, reduced maltose syrup, croscarmellose sodium, brilliant blue

Precautions

When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur)

- This product should not be taken in the following persons
Children under 15 years old.
- Do not use any of the following medications while taking this drug
Other medicine for motion sickness, cold medicine, antipyretics/analgesics, sedatives, expectorant, indigestion and heartburn relief, internal medicine containing antihistamines, etc. (such as internal medicine for rhinitis and medicine for allergies)
- After taking this drug, do not drive a car or operate machinery
(symptoms such as sleepiness, blurred vision and abnormal brightness may occur).

Consultation

- The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration
Patients undergoing medical treatment from a physician.
Pregnant women or women suspected of being pregnant.
The elderly.
Patients who have experienced allergic symptoms associated with drugs, etc.
Persons with the following symptoms.

Urination difficulty

Persons diagnosed as having the following.

Glaucoma, heart disease

- If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation

Skin: rash/redness, itching

Neuropsychiatric system: headache

Urinary system: urination difficulty

Other: facial hot flashes, abnormal brightness

- The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid.
Aplastic anemia: symptoms, such as bruises, nose bleeds, bleeding gums, fever, pale appearance of skin and mucosa, fatigue, palpitations, shortness of breath, becoming dizzy from feeling sick and blood in the urine, etc. may occur.

Agranulocytosis: Symptoms, such as acute high fever, chills and sore throat, etc. may occur.

- The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation

Dry mouth, constipation, diarrhea, and drowsiness, blurred vision

Precautions for storage and handling

- Avoid direct sunlight, and store the product in a cool place with little humidity.
- Store the product beyond the reach of children.
- Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)
- After opening the inner bag (Aluminum bag), store it in a place where it will not be exposed to light, such as returning it to a box, in order to prevent discoloration due to light.
- Do not use the product past the expiration date. Even before the expiration date, use the product within 6 months after it is opened. (to assure the quality)

11

[Disclaimer on Multilingual Product Information]

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.