

## NEW TOKUHON Chill

Excellent antiphlogistic analgesic effect  
Comfortable stimulation  
Improves blood circulation

### Indication

Lumbago, contusions, sprains, stiff shoulders, joint pain, muscular pain, muscular fatigue, frostbite

### Dosage and administration

Apply an appropriate dose onto the lesion several times daily.

#### Precaution

- The use of the drug for children should be supervised by a parent. Children under 13 years old should not use this drug because it may cause tingling sensations.
- Attention should be paid so that the drug does not come into contact with the eyes. If the drug comes into contact with the eyes, immediately rinse well with cold or lukewarm water. If the symptoms are severe, seek medical attention from an ophthalmologist.
- The drug is for external use only. Do not take it orally.
- Do not use the drug in 1 hour before taking a bath. If you take a bath immediately after using the drug, you may experience a strong tingling sensation.
- If your skin is sensitive, do not apply the drug to the same site continuously.
- Apply a small amount on the inner arm before use, and confirm that no rash/redness, itching, eruption, swelling, and pain occurred.
- Dry the skin area to which the drug is applied. When the drug is not dried well, the drug may discolor the clothes.
- If you feel a strong tingling sensation, wash gently the affected site with whipped soap, rinse it off, and cool the site well.

### Ingredient

Active ingredients in 100mL

I-Menthol 6.0g

Glycol salicylate 3.0g

Vitamin E acetate 0.5g

Chlorpheniramine Maleate 0.1g

Nonanoic acid vanillylamide 12mg

#### Excipients

Alcohol, propylene glycol

### Precautions

#### When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur)

- Do not use in the following sites  
Periorbital area, mucosa, etc.  
Eczema, contact dermatitis, open wounds.

#### Consultation

- The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before use  
Patients who have experienced allergic symptoms associated with drugs, etc.
- If you experience any symptoms such as rash/redness, itching, eruption, swelling, or pain after use, these may be adverse reactions, so immediately discontinue the use, and consult with your physician, pharmacist, or registered salesperson with this explanatory document

- If symptoms do not resolve after using the drug for 5 or 6 days, stop using this medicine, and show this document to your physician, pharmacist, or registered salesperson for a consultation

### Precautions for storage and handling

- Keep out of direct sunlight, and store the drug in a cool place with the cap closed.
- Do not place the drug close to any source of flame.
- Store the product beyond the reach of children.
- Do not transfer the drug to other containers. (It may lead to misuse or quality deterioration)
- The alcohol contained in the drug may damage plastic products and the paint of furniture and equipment. Be careful not to stain them with the drug solution.
- Do not use the product past the expiration date.

41

#### 【Disclaimer on Multilingual Product Information】

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.